



Lesseur, C., Diergaarde, B., Olshan, A. F., Wünsch-Filho, V., Ness, A. R., Liu, G., Lacko, M., Eluf-Neto, J., Franceschi, S., Laggiou, P., Macfarlane, G. J., Richiardi, L., Boccia, S., Polesel, J., Kjaerheim, K., Zaridze, D., Johansson, M., Menezes, A. M., Curado, M. P., ... Brennan, P. (2016). Genome-wide association analyses identify new susceptibility loci for oral cavity and pharyngeal cancer. *Nature Genetics*, 48(12), 1544–1550. <https://doi.org/10.1038/ng.3685>

Peer reviewed version

Link to published version (if available):
[10.1038/ng.3685](https://doi.org/10.1038/ng.3685)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Nature Publishing Group at <http://doi.org/10.1038/ng.3685>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Genome-wide association study of oral cavity and pharyngeal cancer

Corina Lesseur¹, Brenda Diergaarde², Andrew F Olshan³, Victor Wünsch-Filho⁴, Andrew R Ness⁵, Geoffrey Liu⁶, Martin Lacko⁷, José Eluf-Neto⁸, Silvia Franceschi¹, Pagona Lagiou¹⁰, Gary J Macfarlane¹¹, Lorenzo Richiardi¹², Stefania Boccia¹³, Jerry Polesel¹⁴, Kristina Kjaerheim¹⁵, David Zaridze¹⁶, Mattias Johansson¹, Ana M Menezes¹⁷, Maria Paula Curado¹⁸, Max Robinson¹⁹, Wolfgang Ahrens²⁰, Cristina Canova²¹, Ariana Znaor^{1,22}, Xavier Castellsagué²³, David I Conway²⁴, Ivana Holcátová²⁵, Dana Mates²⁶, Marta Vilensky²⁷, Claire Healy²⁸, Neonila Szeszenia-Dąbrowska²⁹, Eleonóra Fabiánová³⁰, Jolanta Lissowska³¹, Jennifer R Grandis³², Mark C Weissler³³, Eloiza H Tajara³⁴, Fabio D Nunes³⁵, Marcos B de Carvalho³⁶, Steve Thomas³⁷, Rayjean J Hung³⁸, Wilbert H M Peters³⁹, Rolando Herrero¹, Gabriella Cadoni⁴⁰, H Bas Bueno-de-Mesquita^{41,42,43}, Steffen Annika⁴⁴, Antonio Agudo⁴⁵, Oxana Shan'gina¹⁶, Xiangjun Xiao⁴⁶, Valérie Gaborieau¹, Amélie Chabrier¹, Devasena Anantharaman¹, Paolo Boffetta⁴⁷, Christopher I Amos⁴⁶, James D McKay¹, Paul Brennan^{*1}

Affiliations

¹International Agency for Research on Cancer (IARC/WHO), Lyon, France.

²Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, and University of Pittsburgh Cancer Institute, Pittsburgh, PA, USA.

³Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, and UNC Lineberger Comprehensive Cancer Center, Chapel Hill, NC USA.

⁴Faculdade de Saúde Pública, Universidade de São Paulo, São Paulo, SP, Brazil.

⁵National Institute for Health Research (NIHR) Biomedical Research Unit in Nutrition Diet and Lifestyle at the University Hospitals Bristol NHS Foundation Trust and the University of Bristol and School of Oral and Dental Sciences, University of Bristol, Bristol, UK.

⁶Princess Margaret Cancer Centre, Toronto, Canada.

⁷Department of Otorhinolaryngology, Head and Neck Surgery, Maastricht University Medical Center, Maastricht, Netherlands.

⁸Departamento de Medicina Preventiva, Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil.

¹⁰Department of Hygiene, Epidemiology and Medical Statistics, School of Medicine, National and Kapodistrian University of Athens, Greece.

¹¹School of Medicine, Medical Sciences and Nutrition, University of Aberdeen, Aberdeen, UK.

¹²Department of Medical Sciences, University of Turin, Turin, Italy.

¹³Section of Hygiene, Institute of Public Health, Università Cattolica del Sacro Cuore, Fondazione Policlinico 'Agostino Gemelli', Rome, Italy.

¹⁴Unit of Cancer Epidemiology, CRO Aviano National Cancer Institute, Aviano, Italy

- 38 ¹⁵Cancer Registry of Norway, Oslo, Norway.
- 39 ¹⁶Department of Cancer Epidemiology and Prevention, Institute of Carcinogenesis, N.N. Blokhin
40 Russian Cancer Research Centre of the Russian Ministry of Health, Moscow, Russian
41 Federation.
- 42 ¹⁷Universidade Federal de Pelotas (UFPel), Pelotas, RS, Brazil
- 43 ¹⁸Epidemiology, International Center for Research (CIPE), A.C.Camargo Cancer Center, Sao
44 Paulo, Brazil
- 45 ¹⁹Centre for Oral Health Research, Newcastle University, Newcastle, UK.
- 46 ²⁰Leibniz Institute for Prevention Research and Epidemiology (BIPS), Bremen, Germany.
- 47 ²¹Department of Molecular Medicine, University of Padova, Padova, Italy.
- 48 ²²Croatian National Institute of Public Health, Zagreb, Croatia.
- 49 ²³Institut Català d'Oncologia (ICO)-DIBELL, CIBER-ESP, L'Hospitalet de Llobregat, Catalonia,
50 Spain.
- 51 ²⁴University of Glasgow, Glasgow, and NHS NSS, Edinburgh, UK.
- 52 ²⁵Institute of Hygiene & Epidemiology 1st Faculty of Medicine, Charles University, Prague,
53 Czech Republic.
- 54 ²⁶National Institute of Public Health, Bucharest, Romania.
- 55 ²⁷Instituto de Oncologia "Angel H Roffo", Universidad de Buenos Aires, Buenos Aires,
56 Argentina.
- 57 ²⁸Trinity College School of Dental Science, Dublin, Ireland.
- 58 ²⁹Department of Environmental Epidemiology, Nofer Institute of Occupational Medicine, Lodz,
59 Poland.
- 60 ³⁰Regional Authority of Public Health, Banská Bystrica, Slovakia.
- 61 ³¹The Maria Skłodowska-Curie Memorial Cancer Centre and Institute of Oncology (MCMCC),
62 Warsaw, Poland.
- 63 ³²Department of Otolaryngology Head and Neck Surgery, and Clinical and Translational Science
64 Institute, University of California at San Francisco, San Francisco, CA, USA.
- 65 ³³Department of Otolaryngology/Head and Neck Surgery, University of North Carolina at Chapel
66 Hill, Chapel Hill, NC, USA.
- 67 ³⁴Department of Molecular Biology, School of Medicine of São José do Rio Preto, São José do
68 Rio Preto, SP, Brazil.
- 69 ³⁵Department of Stomatology, School of Dentistry, University of São Paulo, São Paulo, SP,
70 Brazil.
- 71 ³⁶Department of Head and Neck Surgery, Heliópolis Hospital, São Paulo, SP, Brazil
- 72 ³⁷School of Oral and Dental Sciences, University of Bristol, Bristol, UK.
- 73 ³⁸Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, Toronto, Canada.
- 74 ³⁹Department of Gastroenterology, Radboud University Nijmegen Medical Center, Nijmegen,
75 The Netherlands.
- 76 ⁴⁰Institute of Otorhinolaryngology, Università Cattolica del Sacro Cuore, Fondazione Policlinico
77 'Agostino Gemelli', Rome, Italy.
- 78 ⁴¹Department for Determinants of Chronic Diseases (DCD), National Institute for Public Health
79 and the Environment (RIVM), Bilthoven, Netherlands.
- 80 ⁴²Department of Epidemiology and Biostatistics, The School of Public Health, Imperial College
81 London, London, UK.

⁴³Department of Social & Preventive Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia.

⁴⁴German Institute of Human Nutrition in Potsdam-Rehbruecke (DIfE), Nuthetal, Germany.

⁴⁵Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, Catalan Institute of Oncology-IDIBELL, L'Hospitalet de Llobregat, Barcelona, Spain.

⁴⁶Department of Community and Family Medicine Geisel School of Medicine, Dartmouth College, Lebanon, NH, USA.

⁴⁷Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA

§ Xavier Castellsagué recently passed away (6/12/2016)

***Corresponding Author:**

Paul Brennan, Ph.D.
Section of Genetics
International Agency for Research on Cancer (IARC)
Lyon, France
gep@iarc.fr

99 We conducted a genome-wide association study of oral cavity and pharyngeal cancer in
100 6,034 cases and 6,585 controls from Europe, North America and South America. We
101 evaluated >7 million SNPs with oral and pharyngeal cancer risk and detected 8 loci at
102 $P < 5 \times 10^{-8}$, 7 of which are novel for these cancer sites. Overall, oral and pharyngeal
103 cancers were associated with novel loci at 6p21.32 (**rs3828805**, *HLA-DQB1*), 10q26.13
104 (**rs201982221**, *LHHP*) and 11p15.4 (**rs1453414**, *OR52N2/TRIM5*). Oral cancer was
105 associated with two new regions 2p23.3 (**rs6547741**, *GPN1*) and 9q34.12 (**rs928674**,
106 *LAMC3*), and with known cancer susceptibility loci 9p21.3 (**rs8181047**, *CDKN2B-AS1*)
107 and 5p15.33 (**rs10462706**, *CLPTM1L*). Novel oropharyngeal cancer associations were
108 limited to the human leukocyte antigen (HLA) region at 6p21.32 **and** HLA classical allele
109 imputation revealed a protective association with the class II haplotype *DRB1*1301-*
110 *DQA1*0103-DQB1*0603* (odds ratio (OR)=0.59, $P=2.7 \times 10^{-9}$). Stratified analyses on a
111 subgroup of oropharyngeal cases with human papillomavirus (HPV) infection status
112 indicated that this association was considerably stronger in HPV-positive (OR=0.23,
113 $P=1.6 \times 10^{-6}$) compared to HPV-negative cancers (OR=0.75, $P=0.16$).

Cancers of the oral cavity (OC) and oropharynx (OPC) are predominantly caused by tobacco and alcohol use, although oral infection with HPV, particularly HPV16, is an increasingly important cause of OPC¹, especially in the US and northern Europe^{1,2}. The proportion of HPV-related OPCs varies widely and is estimated to be approximately 60% in the US, 30% in Europe and lower in South America²⁻⁵. Genetic factors have also been implicated in OC and OPC susceptibility, especially polymorphisms within alcohol-related genes including alcohol-dehydrogenase 1B (*ADH1B*) and *ADH7*^{6,7}. In order to identify additional susceptibility loci, 13,107 individuals from 12 epidemiological studies (**Supplementary Table 1**) were genotyped using the Illumina OncoArray and after stringent quality-control steps (**Supplementary Table 2, Online Methods**) 6,034 cases and 6,585 cancer-free controls remained for analyses (**Table 1**). We next performed genome-wide imputation using the Haplotype Reference Consortium panel⁸ and obtained approximately 7 million high-quality imputed variants (**Supplementary Fig. 1**). Given the ethnic diversity of our study, we evaluated associations within continent (Europe, North and South America) using multivariate unconditional logistic regressions under a log-additive genetic model adjusted for age, sex and regional eigenvectors. Results by continent were combined using fixed-effect meta-analyses to derive associations for overall OC and pharynx cancer (oral, oropharynx, hypopharynx and overlapping cancers; n=6,034), as well as site-specific OC (n=2,990) and OPC (n=2,641). Although, several ethnic groups are present in the study, supervised ancestry analyses indicated that >90% of participants were predominantly of Caucasian (>70%CEU) ancestry, although some population admixture was observed in South America (**Supplementary Table 3**).

GWA meta-analyses of overall and site-specific cancers identified 9 regions at genome-wide significance ($P < 5 \times 10^{-8}$) (**Fig.1**). Quantile-quantile (Q-Q) plots of observed and expected P -values showed moderate genomic inflation (λ) for the 3 meta-analyses (λ range=1.04-1.06, **Supplementary Fig. 2-3**). Since λ increases with sample size, we scaled it to 1000 cases and controls resulting in ameliorated inflation (λ_{1000} range=1.009-1.01)⁹. Overall OC and pharynx cancer were associated with rs79767424 (5p14.3), rs1229984 (4q23), rs201982221 (10q26.13), rs1453414 (11p15.4) and 123 SNPs at 6p21.32 (**Supplementary Table 4**). Twenty-six variants were associated ($P < 5 \times 10^{-8}$) with OC (**Supplementary Table 5**), 4 of which mapped to 2p23.3, 1 to 4q23, 3 to 9q34.12, 13 to 5p15.33 and 5 to 9p21.3. For OPC, novel significant variants were located at 6p21.32 (62 SNPs, **Supplementary Table 6**). Suggestive susceptibility variants ($P < 5 \times 10^{-7}$, **Supplementary Tables 7-9**) were associated with OC at 4 additional loci: 6p21.33, 6p21.32, 15q21.2, 15q26.2 and, with OPC at 2q36.1. Other genomic locations outside the HLA region showed promising associations ($P < 5 \times 10^{-6}$) with OPC (**Supplementary Table 10**). For susceptibility loci at $P < 5 \times 10^{-8}$, functional annotation of regulatory features with ENCODE and eQTL information, if available, are summarized in **Supplementary Table 11 and 12**. Given the geographical heterogeneity of our population, we performed sensitivity analyses after excluding individuals with <70% CEU ancestry and these showed similar results (**Supplementary Table 13**). To validate array genotypes and imputed dosages, we directly genotyped by a different platform (TaqMan) at least one variant within each locus ($P = 5 \times 10^{-7}$) in a subset of approximately 700 individuals. Concordance between genotyped/imputed genotypes and TaqMan results was >97% for all regions with the exception of

rs2398180, an imputed variant which had a concordance of 94% (**Supplementary Table 14**). For 2 rare variants, rs201982221 (10q26.13) and rs7976742 (5p14.3), TaqMan assays could not be designed and we used Sanger sequencing for validation. We were able to validate the rs201982221 deletion (**Supplementary Fig. 4**), but rs7976742 did not validate (**Online Methods**). The lead variant at each validated loci ($P < 5 \times 10^{-8}$) for overall and site-specific analyses are shown in **Table 2**.

The rs1229984 (4q23, *ADH1B*) association has been previously reported as a susceptibility locus for OC and OPC⁶, and similar to previous findings this variant showed heterogeneity by region, smoking and alcohol drinking status^{10,11} (**Fig. 2a**). Three other 4q23 SNPs reached $P < 5 \times 10^{-8}$, although conditional analyses indicated these are not independent signals (**Supplementary Table 15**). The rs1573496 (*ADH7*) variant reported to be strongly associated with OC and OPC in the previous upper aerodigestive tract cancer GWAS⁷ was only moderately associated here (**Supplementary Table 16**). In the overall OC and pharynx cancer analysis, we identified rs201982221 at 10q26.13 (OR=1.67, $P=1.58 \times 10^{-9}$), that was also separately associated with OC (OR=1.71, $P=1.04 \times 10^{-7}$) and OPC (OR=1.70, $P=7.9 \times 10^{-7}$) (**Fig. 2b**). rs201982221 is located within the *LHPP* gene in a region with reported regulatory features (**Supplementary Table 11**). However, it is a rare intronic deletion in an area of low linkage disequilibrium (LD) (**Supplementary Fig. 5**), and thus warrants further validation in a different population. rs1453414, the lead signal at 11p15.4 (**Supplementary Table 17**), is an intronic variant that showed a borderline association in the overall (OR=1.19, $P=4.78 \times 10^{-8}$) and site-specific analyses [OC (OR=1.19, $P=1.65 \times 10^{-5}$) and OPC (OR=1.22, $P=4.26 \times 10^{-6}$)] (**Fig. 2c**, **Supplemental Fig. 6**).

rs1453414 is upstream of *OR52N2*, an olfactory receptor, and within *TRIM5*, an E3-ubiquitin ligase, and is an eQTL for these genes in brain tissue¹² (**Supplementary Table 12**).

At 2p23.3, 4 SNPs showed evidence ($P < 5 \times 10^{-8}$) for an association with OC, and in conditional analyses did not appear to be independent (**Supplementary Table 18**). These signals map to a high LD area that includes *C2orf16*, *ZNF512*, *CCDC121* and *GPN1* (**Supplementary Fig. 7**). The lead SNP, rs6547741, was associated with OC but not with OPC, and maps to an intron of *GPN1*, a GTPase involved in RNA polymerase II transport and DNA repair¹³. Associations between rs6547741 and OC were homogenous across other stratified analyses by region, sex, smoking and drinking status (**Fig. 3a**).

Variation within 5p15.33 was also exclusively associated with OC (OPC, rs10462706, $P = 0.47$). The top signal, rs10462706, was associated with decreased OC risk ($OR = 0.74$, $P = 5.54 \times 10^{-10}$) and is in low LD ($r^2 = 0.15$, **Supplementary Fig. 8**) with the second strongest signal rs467095. These two variants are 7kb apart and map to intron 13 of *CLPTM1L* and in stratified analysis showed stronger effects in never smokers ($P_{het} = 0.07$ and $P_{het} = 0.0028$, respectively) and never drinkers ($P_{het} = 0.01$ and $P_{het} = 0.0025$, respectively) (**Fig. 3b**, **Supplementary Fig. 9**). Conditional analyses showed that these SNPs are not completely independent (**Supplementary Table 19**). *TERT* and *CLPTM1L* encode the telomerase reverse transcriptase (*TERT*) and the cleft-lip and palate-associated transmembrane 1-like protein (*CLPTM1L*), respectively. Notably, rs467095 is an esophageal *TERT* eQTL¹⁴ (**Supplementary Table 12**) and is in high LD with rs401681 ($OR = 1.18$, $P = 2.1 \times 10^{-7}$, $r^2 = 0.94$) a widely studied SNP associated

with risk of several cancers including: lung^{15,16}, bladder, prostate, cervical, melanoma¹⁷, basal cell¹⁸, esophageal¹⁹, pancreatic²⁰ and nasopharyngeal²¹. Multiple 5p15.33 variants have been reported to independently influence cancer risk in both an increasing and decreasing fashion. Interestingly, rs401681[A] was associated with an increased OC risk similar to previous melanoma associations, and in an opposite direction to previous lung cancer results¹⁷.

Several variants within the *CDKN2A–CDKN2B* locus (9p21.3) were found to be associated with OC. The lead SNP, rs8181047, is an intronic variant within the *CDKN2B1* antisense RNA 1 (*CDKN2B-AS1*) (**Fig. 3c**). rs8181047 is in LD ($r^2_{\text{range}}=0.6-0.8$) with 4 other 9p21.3 variants strongly associated with OC (**Supplementary Fig. 10a**) that in conditional analyses did not show independent associations (**Supplementary Table 20**). The *CDKN2A–CDKN2B* locus contains genes involved in cell-cycle regulation and senescence and has been associated with multiple malignancies including melanoma²², glioma²³, basal cell¹⁸, breast²⁴, lung²⁵, nasopharyngeal²⁶ and esophageal cancer²⁷. Notably, *CDKN2A* is frequently mutated in HPV-negative head and neck cancers²⁸.

The OC associated variants at 9q34.12 mapped to an intron of *LAMC3*, a laminin involved in cortical development²⁹. rs928674, the peak signal, showed consistent effects across strata and a weaker association with OPC ($P=0.003$) (**Fig. 3d**). rs928674 is in high LD with 3 other robustly associated 9q34.12 SNPs ($r^2_{\text{range}}=0.82-0.96$, **Supplemental Fig. 10b**, **Supplementary Table 21**) and is an esophageal mucosa cis-eQTL for a downstream gene *AIF1L* (Allograft Inflammatory Factor 1-Like)¹⁴.

The most prominent finding in the overall and OPC meta-analyses was a large association signal at 6p21.32 within the HLA class II region. The lead variant in both analyses, rs3828805, maps 1.7kb 5' of *HLA-DQB1* (**Fig. 2d** and **Supplementary Fig. 11**) and similar to other 6p21.32 variants (**Supplementary Table 4 and 6**), showed heterogeneity by geographical region ($P_{\text{het}}=0.007$) with no effect in South America ($P=0.62$). Association analyses of 6p21.32 variants ($P<5\times10^{-8}$) conditioned on rs3828805 did not reveal multiple independent signals (**Supplementary Table 22**), suggesting a common haplotype. To further investigate HLA associations, we imputed classical alleles in 11,436 individuals (>70% Caucasian ancestry) (**Online Methods**). Three classical HLA alleles *DRB1*1301*, *DQA1*0103* and *DQB1*0603* reached $P<5\times10^{-8}$ in the overall analysis and were also strongly associated with OPC (**Supplementary Table 23**). These alleles are in high LD ($r^2>0.9$) and are part of the HLA class II haplotype, *DRB1*1301-DQA1*0103-DQB1*0603*, which is common in Europeans and previously reported to be associated with decreased cervical cancer risk³⁰. *DRB1*1301-DQA1*0103-DQB1*0603* was strongly associated with reduced OPC risk (OR=0.59, $P=2.7\times10^{-9}$) and more weakly with OC risk (OR=0.75, $P=1.7\times10^{-4}$). Further conditional analysis on this haplotype and 6p21.32 variants did not reveal evidence of additional independent effects (**Supplementary Table 24**). Given the importance of HPV infection in the etiology of cervical and oropharyngeal cancer³¹, we conducted post-hoc analyses to examine the effect of *DRB1*1301-DQA1*0103-DQB1*0603* in a subset of 576 cases with available HPV-status and 3662 controls. *DRB1*1301-DQA1*0103-DQB1*0603* was associated with a strong reduced risk of HPV-positive OPC (OR=0.23, $P=1\times10^{-6}$, n=336) with no significant association in 240 OPC HPV-negative cases (OR=0.75, $P=0.16$)

(**Table 3**). These results indicate that the class II HLA region is implicated in at least two HPV-driven cancers, namely HPV-positive OPC and cervical cancer. The lack of an association between 6p21.3 SNPs and OPC risk in South America could relate to previous findings that less than 10% of OPC are HPV-positive in this region^{4,5}. Moreover, a weaker association with OC could be due to a smaller proportion of these cases being HPV-positive, as well as possibly some misclassified OPC cases, especially for base of the tongue tumors. Further evaluation of the extent and specificity of this HLA effect in HPV-associated cancers is important given the strength of the observed association. This may help elucidate why some individuals are at higher risk of HPV-positive OPC after HPV infection and may also have implications for cancer immunotherapies targeting the HLA class II antigen presentation pathway³².

In summary, we identified 7 oral and pharyngeal cancer susceptibility loci including a strong HLA signal narrowed to a class II haplotype. Future replication of these findings in an independent population is warranted as well as fine-mapping and functional studies necessary to establish the biological framework underneath these associations.

267 **URLs.** R, <http://www.r-project.org/>; PLINK, <https://www.cog-genomics.org/plink2>;
268 University of Michigan Imputation Server, <https://imputationserver.sph.umich.edu/>;
269 Haplotype Reference Consortium, <http://www.haplotype-reference-consortium.org/>;
270 SNP2HLA, <https://www.broadinstitute.org/mpg/snp2hla/>; HaploReg v4.1,
271 <http://www.broadinstitute.org/mammals/haploreg/haploreg.php>; GTEx Portal,
272 <http://www.gtexportal.org/home/>; LocusZoom,
273 <http://locuszoom.sph.umich.edu/locuszoom/>; metafor R package [http://www.metafor-](http://www.metafor-project.org/doku.php)
274 [project.org/doku.php](http://www.metafor-project.org/doku.php); LDlink, <http://analysistools.nci.nih.gov/LDlink/>; INHANCE
275 Consortium, <http://www.inhance.utah.edu/>; OncoArray Network,
276 <http://epi.grants.cancer.gov/oncoarray/>; GAME-ON,
277 <http://epi.grants.cancer.gov/gameon/>; GENCAPO, <http://www.gencapo.famerp.br>

278

279 **Acknowledgments**

280 We thank Dr. Leticia Fernandez (Instituto Nacional de Oncologia y Radiobiologia, La
281 Habana, Cuba) for her contribution to the IARC ORC multicenter study. We are also
282 grateful to Sergio Koifman (Escola Nacional de Saúde Pública, Rio de Janeiro, Brazil)
283 for his contribution to the IARC Latin America multicenter study (Sergio Koifman passed
284 away in May 2014). We acknowledge all of the participants involved in this research and
285 the funders and support. Genotyping performed at the Center for Inherited Disease
286 Research (CIDR) was funded through the U.S. National Institute of Dental and
287 Craniofacial Research (NIDCR) grant 1X01HG007780-0. Genotyping for shared
288 controls with the Lung OncoArray initiative was funded through the grant
289 X01HG007492-0. Corina Lesseur undertook this work during the tenure of a
290 Postdoctoral Fellowship awarded by the International Agency for Research on Cancer.
291 The funders did not participate in the study design, data collection and analysis,
292 decision to publish, or preparation of the manuscript.

293 The University of Pittsburgh head and neck cancer case-control study is supported by
294 National Institutes of Health grants P50 CA097190 and P30 CA047904. The Carolina
295 Head and Neck Cancer Study (CHANCE) was supported by the National Cancer
296 Institute (R01-CA90731). The Head and Neck Genome Project (GENCAPO) was
297 supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP)
298 (Grant numbers 04/12054-9 and 10/51168-0). The authors thank all the members of the
299 GENCAPO team. The HN5000 study was funded by the National Institute for Health
300 Research (NIHR) under its Programme Grants for Applied Research scheme (RP-PG-
301 0707-10034), the views expressed in this publication are those of the author(s) and not
302 necessarily those of the NHS, the NIHR or the Department of Health. The Toronto study
303 was funded by the Canadian Cancer Society Research Institute (020214) and the
304 National Cancer Institute (U19 CA148127) and the Cancer Care Ontario Research
305 Chair. The alcohol-related cancers and genetic susceptibility study in Europe
306 (ARCAGE) was funded by the European Commission's 5th Framework Program (QLK1-
307 2001-00182), the Italian Association for Cancer Research, Compagnia di San
308 Paolo/FIRMS, Region Piemonte, and Padova University (CPDA057222).The Rome
309 Study was supported by the Associazione Italiana per la Ricerca sul Cancro (AIRC) IG
310 2011 10491 and IG2013 14220 to SB, and Fondazione Veronesi to SB. The IARC Latin
311 American study was funded by the European Commission INCO-DC programme (IC18-
312 CT97-0222), with additional funding from Fondo para la Investigacion Cientifica y
313 Tecnologica (Argentina) and the Fundação de Amparo à Pesquisa do Estado de São
314 Paulo (01/01768-2). The IARC Central Europe study was supported by European
315 Commission's INCO-COPERNICUS Program (IC15-CT98-0332), NIH/National Cancer
316 Institute grant CA92039, and the World Cancer Research Foundation grant WCRF
317 99A28.The IARC Oral Cancer Multicenter study was funded by: grant S06 96 202489
318 05F02 from Europe against Cancer; Grants FIS 97/0024, FIS 97/0662, and BAE
319 01/5013 from Fondo de Investigaciones Sanitarias, Spain; UICC Yamagiwa-Yoshida
320 Memorial International Cancer Study; National Cancer Institute of Canada; Italian
321 Association for Research on Cancer; and the Pan American Health Organization. The
322 coordination of EPIC study is financially supported by the European Commission (DG
323 SANCO) and the International Agency for Research on Cancer.

Competing financial interests: The authors declare no competing financial interests.

Author contributions

P.Brennan and J.D.M. conceived and designed the project. C.L. undertook data harmonization, genotypes quality control, GWAS analysis, imputation and meta-analyses. X.X. performed genotype calling. V.G. and A.C. organized and supervised sample selection and DNA shipments at IARC. A.C. performed replication TaqMan genotyping. C.L. and V.G. analyzed data from replication genotyping. C.L. and P.Brennan drafted the first version of the manuscript. B.D., A.F.O, V.W.-F., A.R.N, G.L., M.L., J.E.-N., S.F., P.L., G.J.M, L.R., S.B., J.P., K.K., D.Z., M.J., A.M.M., M.P.C., M.R., W.A., C.C., A.Z., X.C., D.I.C, I.H., D.M., M.V., C.M.H., N.S.-D., E.F., J.L., J.R.G, M.C.W., E.H.T, F.D.N, M.B.C., S.T., R.J.H., W.H.M.P., R.H., G.C., A.S., A.A., O.S., H.B.B.Dm, P.Boffetta and D.A., contributed with reagent/samples/material and reviewed/approved the final manuscript. J.D.M. and C.I.A designed and coordinated the Lung Cancer OncoArray. P.Brennan obtained funding for the project, provided overall supervision and management.

REFERENCES

1. Gillison, M.L., Chaturvedi, A.K., Anderson, W.F. & Fakhry, C. Epidemiology of Human Papillomavirus-Positive Head and Neck Squamous Cell Carcinoma. *J Clin Oncol* **33**, 3235-42 (2015).
2. Chaturvedi, A.K. *et al.* Human papillomavirus and rising oropharyngeal cancer incidence in the United States. *Journal of Clinical Oncology* **29**, 4294-4301 (2011).
3. Kreimer, A.R., Clifford, G.M., Boyle, P. & Franceschi, S. Human papillomavirus types in head and neck squamous cell carcinomas worldwide: a systematic review. *Cancer Epidemiol Biomarkers Prev* **14**, 467-75 (2005).
4. Ribeiro, K.B. *et al.* Low human papillomavirus prevalence in head and neck cancer: results from two large case-control studies in high-incidence regions. *Int J Epidemiol* **40**, 489-502 (2011).
5. Lopez, R.V. *et al.* Human papillomavirus (HPV) 16 and the prognosis of head and neck cancer in a geographical region with a low prevalence of HPV infection. *Cancer Causes Control* **25**, 461-71 (2014).
6. Hashibe, M. *et al.* Multiple ADH genes are associated with upper aerodigestive cancers. *Nature genetics* **40**, 707-709 (2008).
7. McKay, J.D. *et al.* A genome-wide association study of upper aerodigestive tract cancers conducted within the INHANCE consortium. *PLoS genetics* **7**, e1001333 (2011).
8. McCarthy, S. *et al.* A reference panel of 64,976 haplotypes for genotype imputation. *bioRxiv*, 035170 (2015).
9. de Bakker, P.I. *et al.* Practical aspects of imputation-driven meta-analysis of genome-wide association studies. *Hum Mol Genet* **17**, R122-8 (2008).
10. Hashibe, M. *et al.* Evidence for an important role of alcohol- and aldehyde-metabolizing genes in cancers of the upper aerodigestive tract. *Cancer Epidemiol Biomarkers Prev* **15**, 696-703 (2006).
11. Chang, J.S., Straif, K. & Guha, N. The role of alcohol dehydrogenase genes in head and neck cancers: a systematic review and meta-analysis of ADH1B and ADH1C. *Mutagenesis* **27**, 275-86 (2012).

- 371 12. Ramasamy, A. *et al.* Genetic variability in the regulation of gene expression in
372 ten regions of the human brain. *Nat Neurosci* **17**, 1418-28 (2014).
- 373 13. Carre, C. & Shiekhhattar, R. Human GTPases associate with RNA polymerase II
374 to mediate its nuclear import. *Mol Cell Biol* **31**, 3953-62 (2011).
- 375 14. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis:
376 multitissue gene regulation in humans. *Science* **348**, 648-60 (2015).
- 377 15. McKay, J.D. *et al.* Lung cancer susceptibility locus at 5p15.33. *Nat Genet* **40**,
378 1404-6 (2008).
- 379 16. Wang, Y. *et al.* Common 5p15.33 and 6p21.33 variants influence lung cancer
380 risk. *Nat Genet* **40**, 1407-9 (2008).
- 381 17. Rafnar, T. *et al.* Sequence variants at the TERT-CLPTM1L locus associate with
382 many cancer types. *Nat Genet* **41**, 221-7 (2009).
- 383 18. Stacey, S.N. *et al.* New common variants affecting susceptibility to basal cell
384 carcinoma. *Nat Genet* **41**, 909-14 (2009).
- 385 19. Yin, J. *et al.* TERT-CLPTM1L Rs401681 C>T polymorphism was associated with
386 a decreased risk of esophageal cancer in a Chinese population. *PLoS One* **9**,
387 e100667 (2014).
- 388 20. Petersen, G.M. *et al.* A genome-wide association study identifies pancreatic
389 cancer susceptibility loci on chromosomes 13q22.1, 1q32.1 and 5p15.33. *Nat*
390 *Genet* **42**, 224-8 (2010).
- 391 21. Bei, J.X. *et al.* A GWAS Meta-analysis and Replication Study Identifies a Novel
392 Locus within CLPTM1L/TERT Associated with Nasopharyngeal Carcinoma in
393 Individuals of Chinese Ancestry. *Cancer Epidemiol Biomarkers Prev* **25**, 188-92
394 (2016).
- 395 22. Law, M.H. *et al.* Genome-wide meta-analysis identifies five new susceptibility loci
396 for cutaneous malignant melanoma. *Nat Genet* **47**, 987-95 (2015).
- 397 23. Shete, S. *et al.* Genome-wide association study identifies five susceptibility loci
398 for glioma. *Nat Genet* **41**, 899-904 (2009).
- 399 24. Turnbull, C. *et al.* Genome-wide association study identifies five new breast
400 cancer susceptibility loci. *Nat Genet* **42**, 504-7 (2010).

- 401 25. Timofeeva, M.N. *et al.* Influence of common genetic variation on lung cancer risk:
402 meta-analysis of 14 900 cases and 29 485 controls. *Hum Mol Genet* **21**, 4980-95
403 (2012).
- 404 26. Bei, J.X. *et al.* A genome-wide association study of nasopharyngeal carcinoma
405 identifies three new susceptibility loci. *Nat Genet* **42**, 599-603 (2010).
- 406 27. Wu, C. *et al.* Joint analysis of three genome-wide association studies of
407 esophageal squamous cell carcinoma in Chinese populations. *Nat Genet* **46**,
408 1001-6 (2014).
- 409 28. Comprehensive genomic characterization of head and neck squamous cell
410 carcinomas. *Nature* **517**, 576-82 (2015).
- 411 29. Barak, T. *et al.* Recessive LAMC3 mutations cause malformations of occipital
412 cortical development. *Nat Genet* **43**, 590-4 (2011).
- 413 30. Chen, D. *et al.* Genome-wide association study of susceptibility loci for cervical
414 cancer. *J Natl Cancer Inst* **105**, 624-33 (2013).
- 415 31. Bouvard, V. *et al.* A review of human carcinogens—Part B: biological agents. *The*
416 *lancet oncology* **10**, 321-322 (2009).
- 417 32. Thibodeau, J., Bourgeois-Daigneault, M.C. & Lapointe, R. Targeting the MHC
418 Class II antigen presentation pathway in cancer immunotherapy.
419 *Oncoimmunology* **1**, 908-916 (2012).
- 420

421 **Table 1**
 422 Epidemiological and clinical characteristics of cases and controls.

	Cases		Controls	
	N	%	N	%
Total	6034		6585	
Tumor site				
Oral cavity	2990	49.55		
Oropharynx	2641	43.77		
Hypopharynx	305	5.05		
Overlapping	73	1.21		
Other	25	0.41		
Geographic Region				
Europe	2499	41.42	2928	44.46
North America	2549	42.24	2522	38.30
South America	986	16.34	1135	17.24
Sex				
Male	4527	75.02	4325	65.68
Female	1507	24.98	2260	34.32
Age, years				
=<50	1315	21.79	1355	20.58
50-<60	2006	33.24	1954	29.67
60-<70	1748	28.97	1983	30.11
>=70	964	15.98	1293	19.64
Unknown	1	0.02	0	0.00
Smoking Status				
Never	1057	17.52	2508	38.09
Former	1792	29.70	2263	34.37
Current	2623	43.47	1466	22.26
Unknown	562	9.31	348	5.28
Drinking Status				
Never	820	13.59	1199	18.21
Ever	4840	80.21	4840	73.50
Unknown	374	6.20	546	8.29

Table 2

Lead genome-wide significant SNP per validated locus from the regional meta-analyses of OC^a and pharynx cancer combined, as well as OC^a and OPC^a separately.

Region	SNP	chr:pos ^b	Gene	EA/ OA ^c	Info (Rsqr) ^d	AF ^e case/ control	OR	P	P _{het}
Oral and pharyngeal cancer									
4q23	rs1229984	4:100239319	<i>ADH1B</i>	A/G	Geno	0.03/0.06	0.56	2.29x10 ⁻¹⁵	0.002
6p21.32	rs3828805	6:32636120	<i>HLA-DQB1</i>	C/T	0.88	0.75/0.72	1.28	3.35x10 ⁻¹³	0.007
10q26.13	rs20198222 1	10:126157446	<i>LHPP</i>	D/I	Geno	0.03/0.02	1.67	1.58x10 ⁻⁹	0.50
11p15.4	rs1453414	11:5829084	<i>OR52N2/TRIM5</i>	C/A	Geno	0.23/0.20	1.19	4.78x10 ⁻⁸	0.55
Oral cancer									
2p23.3	rs6547741	2:27855924	<i>GPN1</i>	A/G	0.98	0.50/0.54	0.83	3.97x10 ⁻⁸	0.34
4q23	rs1229984	4:100239319	<i>ADH1B</i>	A/G	Geno	0.03/0.06	0.57	1.09x10 ⁻⁹	0.02
5p15.33	rs10462706	5:1343794	<i>CLPTM1L</i>	T/C	0.97	0.12/0.15	0.74	5.54x10 ⁻¹⁰	0.84
9p21.3	rs8181047	9:22064465	<i>CDKN2B-AS1</i>	A/G	Geno	0.29/0.24	1.24	3.80x10 ⁻⁹	0.37
9q34.12	rs928674	9:133952024	<i>LAMC3</i>	G/A	0.89	0.14/0.12	1.33	2.09x10 ⁻⁸	0.88
Oropharyngeal cancer									
4q23	rs1229984	4:100239319	<i>ADH1B</i>	A/G	Geno	0.02/0.06	0.55	8.53x10 ⁻⁹	0.05
6p21.32	rs3828805	6:32636120	<i>HLA-DQB1</i>	C/T	0.88	0.75/0.7	1.37	2.21x10 ⁻¹²	0.07

^a OC=oral cancer, OPC=oropharyngeal cancer; ^b SNP position according to NCBI genome build 37 (Hg19); ^c EA=Effect allele; OA=other allele; ^d Geno=genotyped, SNP, INFO, R² is the average across imputation batches; ^e AF= allele frequency of the effect allele

Figure 1

Genome-wide associations meta-analyses results. The red line represents $P=5 \times 10^{-8}$. The y-axis represents the $-\log_{10} P$ -values. (a) Overall OC and pharyngeal cancer 6,034 cases and 6585 controls (b) Oral cancer analysis with 2,990 cases and 6,585 controls (c) Oropharyngeal cancer analysis with 2,641 cases and 6,585 controls. Loci with GWA significant SNPs and technically validated are tagged with genomic location.




Figure 2

Forest plots of odds ratios for the lead SNP at each genome-wide significant loci in the overall oral and pharyngeal cancer meta-analysis. (a)4q23, rs1229984 (b)10q26.13, rs20198222 (c)11p15.4 rs1453414. (d)6p21.32 rs3828805. EAF=effect allele frequency in 6585 controls. Effect allele in square brackets. OC=oral cancer; OPC=oropharynx cancer. **Plot data in Supplementary Table 25.**







Figure 3

Forest plots of odds ratios for the lead SNP at each genome-wide significant loci in the oral cancer (OC) meta-analysis. (a)2p23.3 rs6547741 (b)5p15.33, rs10462706 (c)9p21.3, rs8181047 (d)9q34.12, rs928674. EAF=effect allele frequency in 6585 controls. Effect allele in square brackets. Overall=oral and pharyngeal cancer; OPC=oropharynx cancer. **Plot data in Supplementary Table 26.**




Table 3Associations of *DRB1*1301-DQA1*0103-DQB1*0603* haplotype in individuals with >70% Caucasian ancestry

	haplotype ^a	case/	HF ^b	HF ^b	OR	P	Meta-analysis ^c	
	case/control	control	case	control			OR	P
Oral and pharynx cancer							0.68	3.32x10 ⁻¹⁰
Europe	207/422	2497/2928	0.04	0.07	0.60	4.04x10 ⁻⁸		
North America	207/276	2342/2329	0.04	0.06	0.74	1.68x10 ⁻³		
South America	74/101	647/727	0.06	0.07	0.86	0.35		
Oral cancer							0.75	1.72x10 ⁻⁴
Europe	106/422	1231/2928	0.04	0.07	0.60	1.52x10 ⁻⁵		
North America	128/276	1135/2329	0.06	0.06	0.92	4.67x10 ⁻¹		
South America	41/101	351/727	0.06	0.07	0.80	0.26		
Oropharynx cancer							0.59	2.73x10 ⁻⁹
Europe	84/422	1098/2928	0.04	0.07	0.57	2.69x10 ⁻⁵		
North America	72/276	1119/2329	0.03	0.06	0.52	3.49x10 ⁻⁶		
South America	31/101	216/727	0.07	0.07	1.05	0.81		
Oropharynx cancer by HPV status^d								
HPV-positive	11/505	336/3686	0.01	0.07	0.23	1.6x10 ⁻⁶		
HPV-negative	25/505	240/3686	0.05	0.07	0.75	0.16		

^a Number of copies of the haplotype in cases and controls^b Haplotype frequency calculated as total number of copies of haplotype in the population (haplotype copies/2n).^c Fixed-effects meta-analysis of regional associations adjusted for age, sex and eigenvectors.

^dHPV-status available in a subset of cases from ARCAGE, EPIC, CHANCE and Pittsburgh studies